

Hadronic light-by-light scattering contribution to the muon $g - 2^*$

Andreas Nyffeler^a

^aCentre de Physique Théorique, CNRS-Luminy, Case 907
F-13288 Marseille Cedex 9, France
E-mail: nyffeler@cpt.univ-mrs.fr

We briefly review the current status of the hadronic light-by-light scattering correction to the muon $g - 2$. Then we present our semi-analytical evaluation of the pion-pole contribution, using a description of the pion-photon-photon form factor based on large- N_C and short-distance properties of QCD. We also sketch an effective field theory approach to hadronic light-by-light scattering. In view of several still unsolved problems, our conservative estimate for the full hadronic light-by-light scattering contribution is $a_\mu^{\text{LbyL;had}} = +8(4) \times 10^{-10}$.

1. Introduction

The present picture of hadronic light-by-light scattering is shown in Fig. 1 and the corre-




Figure 1. The hadronic light-by-light scattering contribution to the muon $g - 2$.

sponding contributions to a_μ are listed in Table 1, taking into account the corrections made in the two full evaluations [1,2], after we had discovered the sign error in the pion-pole contribution [3,4]. There are three classes of contributions to the

*Talk presented at the 9th International High-Energy Physics Conference in Quantum Chromodynamics (QCD 2002), Montpellier, France, 2–9 July 2002.

Table 1

Contributions to $a_\mu (\times 10^{10})$ according to Fig. 1. The last column gives the result when no form factors are used in the couplings to the photons.

Type	Ref. [1]	Ref. [2]	Ref. [3]	
(b)	-0.5(0.8)	-1.9(1.3)		-4.5
(c)	8.3(0.6)	8.5(1.3)	8.3(1.2)	$+\infty$
f_0, a_1	0.174 ^a	-0.4(0.3)		
(d)	1.0(1.1)	2.1(0.3)		~ 6
Total	9.0(1.5)	8.3(3.2)	8(4) ^b	

^a Only a_1 exchange.

^b Our estimate, using Refs. [1–3].

hadronic four-point function [Fig. 1(a)], which can be understood from an effective field theory (EFT) analysis of hadronic light-by-light scattering [5,4]: (1) a charged pion loop [Fig. 1(b)], where the coupling to photons is dressed by some form factor (ρ -meson exchange, e.g. via vector meson dominance (VMD)), (2) the pseudoscalar pole diagrams [Fig. 1(c)] together with the exchange of heavier resonances ($f_0, a_1, \dots$) and, finally, (3) the irreducible part of the four-point function which was modeled in Refs. [1,2] by a constituent quark loop dressed again with VMD form factors [Fig. 1(d)]. The latter can be viewed as a local contribution $\bar{\psi}\sigma^{\mu\nu}\psi F_{\mu\nu}$ to $g - 2$. The two groups [1,2] used similar, but not identical models which explains the slightly different results for the dressed charged pion and the dressed

constituent quark loop, although their sum seems to cancel to a large extent and the final result is essentially given by the pseudoscalar exchange diagrams. Since the models used in Refs. [1,2] are not fully consistent with QCD, we take the difference of the results as indication of the error coming from the model dependence.

On the other hand, we will show in Section 2 that the pseudoscalar contribution now seems under control, due to our semi-analytical calculation [3], based on a pion-photon-photon form factor $\mathcal{F}_{\pi^0\gamma^*\gamma^*}$ which fulfills the relevant QCD short-distance constraints [6], in contrast to the form factors used in Refs. [1,2]. These findings are also corroborated by an EFT and large- N_C analysis of $a_\mu^{\text{LbyL;had}}$ [4] which allows to calculate the leading and next-to-leading logarithms (Sec. 3).

2. Pion-pole contribution

The contribution from the neutral pion intermediate state is given by a two-loop integral that involves the convolution of two pion-photon-photon transition form factors, see Fig. 1(c). Since no data on the doubly off-shell form factor $\mathcal{F}_{\pi^0\gamma^*\gamma^*}(q_1^2, q_2^2)$ is available, one has to resort to models. In order to proceed with the analytical evaluation of the two-loop integrals, we considered a certain class of form factors which includes the ones based on large- N_C QCD that we studied in Ref. [6]. For comparison, we have also used a vector meson dominance (VMD) and a constant form factor, derived from the Wess-Zumino-Witten (WZW) term. For all these form factors we could perform *all* angular integrations in the two-loop integrals analytically [3].

In large- N_C QCD, the pion-photon-photon form factor is described by a sum over an infinite set of narrow vector resonances, involving arbitrary couplings, although there are constraints at long and short distances. The normalization is given by the WZW term, $\mathcal{F}_{\pi^0\gamma^*\gamma^*}(0, 0) = -N_C/(12\pi^2 F_\pi)$, whereas the OPE tells us that

$$\lim_{\lambda \rightarrow \infty} \mathcal{F}_{\pi^0\gamma^*\gamma^*}(\lambda^2 q^2, (p - \lambda q)^2) = \frac{2}{3} \frac{F_\pi}{q^2} \left\{ \frac{1}{\lambda^2} + \frac{1}{\lambda^3} \frac{q \cdot p}{q^2} + \mathcal{O}\left(\frac{1}{\lambda^4}\right) \right\}. \quad (1)$$

In the following, we consider the form factors that

are obtained by truncation of the infinite sum in large- N_C QCD to one (lowest meson dominance, LMD), and two (LMD+V), vector resonances per channel, respectively:

$$\mathcal{F}_{\pi^0\gamma^*\gamma^*}^{\text{LMD}}(q_1^2, q_2^2) = \frac{F_\pi}{3} \frac{q_1^2 + q_2^2 - c_V}{(q_1^2 - M_V^2)(q_2^2 - M_V^2)}, \quad (2)$$

$$\mathcal{F}_{\pi^0\gamma^*\gamma^*}^{\text{LMD+V}}(q_1^2, q_2^2) = \frac{F_\pi}{3} \left\{ \left(q_1^2 q_2^2 (q_1^2 + q_2^2) + h_1 (q_1^2 + q_2^2)^2 + h_2 q_1^2 q_2^2 + h_5 (q_1^2 + q_2^2) + h_7 \right) / \left((q_1^2 - M_{V_1}^2) \times (q_1^2 - M_{V_2}^2)(q_2^2 - M_{V_1}^2)(q_2^2 - M_{V_2}^2) \right) \right\}, \quad (3)$$

with the constants $c_V = N_C M_V^4 / (4\pi^2 F_\pi^2)$ and $h_7 = -N_C M_{V_1}^4 M_{V_2}^4 / (4\pi^2 F_\pi^2)$. The parameters h_1, h_2 , and h_5 in the LMD+V form factor are not fixed by the normalization and the leading term in the OPE. We have determined these coefficients phenomenologically [6,3]. In particular, $\mathcal{F}_{\pi^0\gamma^*\gamma^*}(-Q^2, 0)$ with one photon on-shell behaves like $1/Q^2$ for large spacelike momenta, $Q^2 = -q^2$. Whereas the LMD form factor does not have such a behavior, it can be reproduced with the LMD+V ansatz, provided that $h_1 = 0$. A fit to the data yields moreover $h_5 = 6.93 \pm 0.26 \text{ GeV}^4$. Analyzing the experimental data for the decay $\pi^0 \rightarrow e^+ e^-$ leads to the loose bound $|h_2| \lesssim 20 \text{ GeV}^2$.

Note that the usual VMD form factor $\mathcal{F}_{\pi^0\gamma^*\gamma^*}^{\text{VMD}}(q_1^2, q_2^2) \sim 1/[(q_1^2 - M_V^2)(q_2^2 - M_V^2)]$ does *not* correctly reproduce the OPE in Eq. (1).

After performing the angular integrations, the pion-exchange contribution to a_μ can be written as a two-dimensional integral representation, where the integration runs over the moduli of the Euclidean momenta

$$a_\mu^{\text{LbyL};\pi^0} = \int_0^\infty dQ_1 \int_0^\infty dQ_2 \times \sum_i w_i(Q_1, Q_2) f_i(Q_1, Q_2), \quad (4)$$

with universal [for the above class of form factors] weight functions w_i (rational functions, square roots and logarithms) [3]. The dependence on the form factors resides in f_i . In this way we could

separate the generic features of the pion-pole contribution from the model dependence. This is not possible anymore in the final analytical result (as a series expansion) for $a_\mu^{\text{LbyL};\pi^0}$ in Ref. [7]. One has to keep in mind that there is an intrinsic uncertainty in the form factor of 10–30 %, furthermore the VMD form factor used in that reference has the wrong high-energy behavior.

The weight functions w_i in the main contribution are positive and peaked around momenta of the order of 0.5 GeV. There is, however, a tail in one of these functions, which produces for the constant WZW form factor a divergence of the form $\ln^2\Lambda$ for some UV-cutoff Λ . Other weight functions have positive and negative contributions in the low-energy region, which lead to a strong cancellation in the corresponding integrals.

In Table 2 we present the numerical results for the different form factors. All form factors lead to very similar results (apart from WZW). Judging from the shape of the weight functions described above, it seems more important to correctly reproduce the slope of the form factor at the origin and the available data at intermediate energies. On the other hand, the asymptotic behavior at large Q_i seems not very relevant. The results for the LMD+V form factor are rather stable under the variation of the parameters, except for h_2 . If all other parameters are kept fixed, our result changes in the range $|h_2| < 20 \text{ GeV}^2$ by $\pm 0.9 \times 10^{-10}$ from the value for $h_2 = 0$.

Thus, with the LMD+V form factor, we get

$$a_\mu^{\text{LbyL};\pi^0} = +5.8 (1.0) \times 10^{-10}, \quad (5)$$

Table 2

Results for $a_\mu^{\text{LbyL};\pi^0}$ for the different form factors. In the WZW model we used a cutoff of 1 GeV in the divergent contribution.

Form factor	$a_\mu^{\text{LbyL};\pi^0} \times 10^{10}$
WZW	12.2
VMD	5.6
LMD	7.3
LMD+V ($h_2 = 0 \text{ GeV}^2$)	5.8

where the error includes the variation of the parameters and the intrinsic model dependence. A similar short-distance analysis in the framework of large- N_C QCD and including quark mass corrections for the form factors for the η and η' was beyond the scope of Ref. [3]. We therefore used VMD form factors fitted to the available data for $\mathcal{F}_{\pi^0\gamma^*\gamma^*}(-Q^2, 0)$ to obtain our final estimate

$$\begin{aligned} a_\mu^{\text{LbyL};\text{PS}} &\equiv a_\mu^{\text{LbyL};\pi^0} + a_\mu^{\text{LbyL};\eta}|_{\text{VMD}} + a_\mu^{\text{LbyL};\eta'}|_{\text{VMD}} \\ &= +8.3 (1.2) \times 10^{-10}. \end{aligned} \quad (6)$$

An error of 15 % for the pseudoscalar pole contribution seems reasonable, since we impose many theoretical constraints from long and short distances on the form factors. Furthermore, we use experimental information whenever available.

3. EFT approach to $a_\mu^{\text{LbyL};\text{had}}$

In Ref. [4] we discussed an EFT approach to hadronic light-by-light scattering based on an effective Lagrangian that describes the physics of the Standard Model well below 1 GeV. It includes photons, light leptons, and the pseudoscalar mesons and obeys chiral symmetry and $U(1)$ gauge invariance.

The leading contribution to $a_\mu^{\text{LbyL};\text{had}}$, of order $\mathcal{O}(p^6)$, is given by a (finite) loop of charged pions with point-like electromagnetic vertices. Since this contribution involves a loop of hadrons, it is subleading in the large- N_C expansion.

At order p^8 and at leading order in N_C , we encounter the divergent pion-pole contribution, diagrams (a) and (b) of Fig. 2, involving two WZW vertices. The diagram (c) is actually finite. The divergences of the triangular subgraphs in the diagrams (a) and (b) are removed by inserting the counterterm χ from the Lagrangian $\mathcal{L}^{(6)} = (\alpha^2/4\pi^2 F_0) \chi \bar{\psi} \gamma_\mu \gamma_5 \psi \partial^\mu \pi^0 + \dots$, see the one-loop diagrams (d) and (e). Finally, there is an overall divergence of the two-loop diagrams (a) and (b) that is removed by a local counterterm, diagram (f). Since the EFT involves such a local contribution, we will not be able to give a precise numerical prediction for $a_\mu^{\text{LbyL};\text{had}}$.

Nevertheless, it is interesting to consider the leading and next-to-leading logarithms that are in addition enhanced by a factor N_C and which can




Figure 2. The graphs contributing to $a_\mu^{\text{LbyL};\pi^0}$ at lowest order in the effective field theory.

be calculated using the renormalization group [4]. The EFT and large- N_C analysis tells us that

$$a_\mu^{\text{LbyL};\text{had}} = \left(\frac{\alpha}{\pi}\right)^3 \left\{ f\left(\frac{m_{\pi^\pm}}{m_\mu}, \frac{m_{K^\pm}}{m_\mu}\right) + N_C \left(\frac{m_\mu^2}{16\pi^2 F_\pi^2} \frac{N_C}{3} \right) \left[\ln^2 \frac{\mu_0}{m_\mu} + c_1 \ln \frac{\mu_0}{m_\mu} + c_0 \right] + \mathcal{O}\left(\frac{m_\mu^2}{\mu_0^2} \times \log^2\right) + \mathcal{O}\left(\frac{m_\mu^4}{\mu_0^4} N_C \times \log\right) \right\}, \quad (7)$$

where $f(m_{\pi^\pm}/m_\mu, m_{K^\pm}/m_\mu) = -0.038$ represents the charged pion and kaon-loop that is formally of order one in the chiral and N_C counting and μ_0 denotes some hadronic scale, e.g. M_ρ . The coefficient $\mathcal{C}$ of the log-square term in the second line is universal and of order N_C , since $F_\pi = \mathcal{O}(\sqrt{N_C})$.

Unfortunately, although the logarithm is sizeable, in $a_\mu^{\text{LbyL};\pi^0}$ there occurs a cancellation between the log-square and the log-term. If we fit our result for the VMD form factor for large M_ρ to an expression as given in Eq. (7), we obtain

$$\begin{aligned} a_{\mu;\text{VMD}}^{\text{LbyL};\pi^0} &\doteq \left(\frac{\alpha}{\pi}\right)^3 \mathcal{C} \left[\ln^2 \frac{M_\rho}{m_\mu} + c_1 \ln \frac{M_\rho}{m_\mu} + c_0 \right] \\ &\stackrel{\text{Fit}}{=} \left(\frac{\alpha}{\pi}\right)^3 \mathcal{C} [3.94 - 3.30 + 1.08] \\ &= [12.3 - 10.3 + 3.4] \times 10^{-10} \\ &= 5.4 \times 10^{-10}, \end{aligned} \quad (8)$$

which is confirmed by the analytical result of Ref. [7] (setting for simplicity $m_{\pi^0} = m_\mu$): $a_{\mu;\text{VMD}}^{\text{LbyL};\pi^0} = [12 - 8.0 + 1.7] \times 10^{-10} = 5.7 \times 10^{-10}$.

This cancellation is now also visible in the revised version of Ref. [8]. In that paper the remaining parts of c_1 have been calculated: $c_1 = -2\chi(\mu_0)/3 + 0.237 = -0.93^{+0.67}_{-0.83}$, with our conventions for χ and $\chi(M_\rho)_{\text{exp}} = 1.75^{+1.25}_{-1.00}$.

4. Conclusions

The pseudoscalar pole contribution $a_\mu^{\text{LbyL};\text{PS}}$ seems to be under control at the 15 % level. Moreover, the EFT and large- N_C analysis shows a systematic approach to $a_\mu^{\text{LbyL};\text{had}}$ and yields the leading and next-to-leading logarithmic terms, enhanced by a factor N_C , although these terms tend to cancel each other. However, there remains the issue of the other contributions, i.e. the dressed charged pion and the dressed constituent quark loop, see Fig. 1 and Table 1, where model calculations lead to slightly different results. Taking these uncertainties into account by adding the errors linearly, my estimate for the full hadronic light-by-light scattering contribution reads:

$$a_\mu^{\text{LbyL};\text{had}} = +8(4) \times 10^{-10}. \quad (9)$$

Acknowledgments

I would like to thank the organizers of QCD 02, in particular S. Narison, for providing such a pleasant atmosphere and the Schweizerischer Nationalfonds for financial support.

REFERENCES

1. M. Hayakawa and T. Kinoshita, Phys. Rev. D **66**, 019902 (E) (2002) and hep-ph/0112102.
2. J. Bijnens, E. Pallante, and J. Prades, Nucl. Phys. **B626**, 410 (2002).
3. M. Knecht and A. Nyffeler, Phys. Rev. D **65**, 073034 (2002).
4. M. Knecht, A. Nyffeler, M. Perrottet, and E. de Rafael, Phys. Rev. Lett. **88**, 071802 (2002).
5. E. de Rafael, Phys. Lett. B **322**, 239 (1994).
6. M. Knecht and A. Nyffeler, Eur. Phys. J. C **21**, 659 (2001).
7. I. Blokland, A. Czarnecki, and K. Melnikov, Phys. Rev. Lett. **88**, 071803 (2002).
8. M. Ramsey-Musolf and M.B. Wise, Phys. Rev. Lett. **89**, 041601 (2002).